

## REFERENCES

1. Guest JF et al . Impacted cerumen: composition,production,epidemiology and management for clients of all ages. *Nurse Pract* 1993 March; 18(3):29
2. Pothier, DD, Hall, C, Gillett, S. A comparison of endoscopic and microscopic removal of wax: a randomised clinical trial. *Clin Otolaryngol* 2006;31:375-80
3. Easton J& Leverton Supporting adults with hearing loss in primary care: new NICE guideline. *BJGP* Nov 2018.
4. Sharp, JF, Wilson, JA, Ross, L, Barr-Hamilton, RM. Ear wax removal: a survey of current practice. *BMJ* 1990;301:1251-3.
5. Addams-Williams, J, Howarth, A, Philipps, JJ. Microsuction aural toilet in ENT outpatients: a questionnaire to evaluate the patient experience. *Eur Arch Otorhinolaryngol* 2010;267:1863-6
6. SJ Prowse, O Mulla . Aural microsuction for wax impaction: survey of efficacy and patient perception. *The Journal of Laryngology & Otology*, 2014 - cambridge.org
7. Snelling, JD, Smithard, A, Waddell, A. Noise levels generated within the external auditory canal during microsuction aural toilet and the effect on hearing: a prospective controlled series. *Clin Otolaryngol* 2009;34:21-5.
8. Burton, MJ, Doree, C. Ear drops for the removal of ear wax. *Cochrane Database Syst Rev* 2009;(2):CD004326
9. Eekhof, JA, De Bock, GH, Le Cessie, S, Springer, MP. A quasi-randomised controlled trial of water as a quick softening agent of persistent earwax in general practice. *Br J Gen Pract* 2001;51:635-7